

Predictability of Particle Trajectories in the Ocean

Tamay M. Ozgokmen, Annalisa Griffa, Arthur J. Mariano
Meteorology and Physical Oceanography Department,
Rosenstiel School of Marine and Atmospheric Science
4600 Rickenbacker Causeway
Miami, Florida 33149

phone: (305) 361 4053 fax: (305) 361 4696 email: tozgokmen@rsmas.miami.edu
Award #: N000149910049

LONG TERM GOALS

The long term goal of this grant is to determine optimal employment and Lagrangian data assimilation strategies for drifting buoys, in order to enhance prediction of particle motion in the ocean with potential applications to ecological, search and rescue, and the floating mine problems.

OBJECTIVES

The specific scientific objective of the work done has been to determine the effectiveness of using in-situ Lagrangian measurements and data assimilation techniques in improving the prediction of particle trajectories. This has been done initially in the context of an ocean model (MICOM) and then using real oceanic drifter data representing both small scale (the Adriatic Sea) and large scale (tropical Pacific Ocean) motion.

APPROACH

The work is based primarily on simple probabilistic particle models and data assimilation strategies. It also involves the use of OGCMs and processing of oceanic data.

WORK COMPLETED

A new method has been developed to address the problem of prediction of Lagrangian trajectories, considering the fact the Lagrangian data are especially suitable for practical applications since they move with the currents and can be easily released in situ. This method employs a simple data assimilation scheme to incorporate information from nearby buoys in order to estimate the turbulent flow characteristics and to improve the prediction of particle trajectories. This method has been initially developed in the context of an OGCM (MICOM), and the results are described in a paper (in press). Since it is portable, and it can easily be applied to real oceanic Lagrangian data as well. The work on a small scale application, using Lagrangian data from the Adriatic Sea is completed and the manuscript is submitted.

RESULTS

The main results, obtained in collaboration with other scientists, can be summarized as follows:

1) The predictability of particle trajectories in oceanic flows is investigated in the context of a primitive-equation, idealized, double-gyre ocean model. The original aspect of this study is the use of

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Predictability of Particle Trajectories in the Ocean				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Miami, Rosenstiel School of Marine and Atmospheric Science, 4600 Rickenbacker Causeway, Miami, FL, 33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

Lagrangian drifter data to improve the accuracy of predicted trajectories. The prediction is performed by assimilating velocity data from the surrounding drifters into a Gauss-Markov model for particle motion. The assimilation is carried out using a simplified Kalman filter.

The performance of the prediction scheme is quantified as a function of a number of factors: (i) dynamically-different flow regimes: interior gyre, western boundary current and midlatitude jet regions; (ii) density of drifter data used in assimilation; and (iii) uncertainties in the knowledge of the mean flow field and the initial conditions. The simulations indicate that predictions improve significantly when the number of data per degrees of freedom, $NR > 1$. Even when the mean flow field and initial turbulent velocities are not known accurately, the information derived from the surrounding drifter data is shown to compensate when $NR > 1$. Theoretical error estimates are derived which are based on the main statistical parameters of the flow field. Theoretical formulae show good agreement with the numerical results and hence they may serve as useful a-priori estimates of Lagrangian prediction error for practical applications.

This work has been performed in collaboration with L. I. Piterbarg (U. Southern California).

2) To test the new prediction scheme with real drifter data, predictability of Lagrangian particle trajectories in the Adriatic Sea (a semi-enclosed sub-basin of the Mediterranean Sea) is investigated using three clusters consisting of 5-7 drifters each over a period of 1-2 weeks. The results of this study (see Fig. 1 for a brief introduction) conducted with real drifter data confirm those with synthetic data that the assimilation of the drifters is efficient for $NR > 1$, while it does not improve the prediction for $NR < 1$. Also, when the mean flow is assumed unknown or an error in the initial position smaller than the Rossby radius of deformation is introduced, predictions are still in good agreement with observations.

This work has been performed in collaboration with P.-M. Poulain (NPS, Monterey).

3) To complement the OCGM and small-scale application (the Adriatic Sea) studies of the new method, we have also started to test it using WOCE drifter data in the tropical Pacific Ocean for large-scale applications. Several clusters of drifters have been tested, with results confirming those of previous studies, hence indicating the validity of our technique for flows for a wide range of scales. Several model improvements have been implemented, most importantly: i) generalization of the Kalman filtering technique for drifter positions and well as velocities; and ii) incorporation of the Coriolis acceleration in the equations of particle motion. This study also emphasizes the difference between estimating (theoretically) the prediction error for a cluster and for a uniformly distributed drifter ensemble (implemented in our previous study outlined in (1)).

This work is being performed in collaboration with L. I. Piterbarg (U. Southern California).




Figure 1: (a) Trajectories of drifter clusters released in the Adriatic Sea (Poulain, 1999). (b) True and simulated trajectories for drifter # 24098 from Castellari et al. (1999). Solid line indicates the true trajectory, simulated trajectory using only the mean flow information is shown with dashed-dotted line, and dashed line depicts the predicted trajectory using our model (11)–(12). Trajectories are marked at 5 day intervals. Similar results have been obtained for the other drifters in this cluster. (c) Absolute dispersion (average distance travelled) of particles, (long dashed line), rms prediction error using only the mean flow (short dashed line) and rms prediction error using the Lagrangian data assimilation model (solid line).

IMPACT/APPLICATIONS

The investigation of the predictability of particle motion is an important area of study, with a number of potential practical applications at very different scales, including searching for persons or valuable objects lost at sea, tracking floating mines, ecological problems such as the spreading of pollutants or fish larvae, and estimating mass transport for climate studies.

TRANSITIONS

Theoretical work is ongoing with our close collaborator L. I. Piterbarg, for the clarification of the difference between the prediction error for a cluster and for a uniformly distributed drifter ensemble, and for the development of appropriate theoretical estimates for clusters.

Improvement of model dynamics is being planned. Considering the fact that for many of the practical applications of the project, the behavior of particles at the surface of the ocean is of importance, the incorporation of wind forcing into the governing equations for particle motion is being planned. A reapplication of the improved model to the Adriatic drifter data set will be conducted using the NORAPS wind data provided by P.-M. Poulain. We anticipate that P.-M. Poulain will be a close collaborator during this study, as he was in our previous Adriatic study.

Ensemble Kalman Filtering techniques are planned to be tested in collaboration with M. Chin (JPL).

Particle predictability study is planned for the drifter data set in the Caribbean Sea, in collaboration with K. Leaman (RSMAS).

RELATED PROJECTS

Lagrangian data analysis in mesoscale prediction and model validation studies. PIs: A. Griffa, T.M. Ozgokmen.

Statistical Problems in Ocean Modeling and Prediction. PI: L.I. Piterbarg.

REFERENCE

Poulain, P.-M., 1999: Drifter observations of surface circulation in the Adriatic Sea between December 1994 and March 1996. *J. Mar. Sys.*, 20, 231-253.

PUBLICATIONS (1998-1999)

Ozgokmen, T.M., A. Griffa, L. Piterbarg, and A. J. Mariano, 1998: Assimilation of Lagrangian data for the prediction of particle motions in a numerical ocean model. *EOS Transactions of the American Geophysical Union*, 79(1), OS36.

Ozgokmen, T.M., 1998: On the predictability of Lagrangian trajectories in the ocean. Conference on Mathematical Problems in Meteorology and Oceanography, Indiana University. Invited talk.

Ozgokmen, T.M., A. Griffa, L. I. Piterbarg, and A. J. Mariano, 1999: Predictability of trajectories in the ocean and Lagrangian data assimilation. International Liege Colloquium on Ocean Hydrodynamics.

Castellari S., A. Griffa, T. M. Ozgokmen, and P.-M. Poulain, 1999: Prediction of particle trajectories in the Adriatic Sea using Lagrangian data assimilation. International Liege Colloquium on Ocean Hydrodynamics.

Ozgokmen, T. M., A. Griffa, L. I. Piterbarg, and A. J. Mariano, 1999: On the predictability of Lagrangian trajectories in the ocean. Journal of Atmospheric and Oceanic Technology, accepted in March 1999.

Castellari S., A. Griffa, T. M. Ozgokmen , and P.-M. Poulain, 1999: Prediction of particle trajectories in the Adriatic Sea using Lagrangian data assimilation. Submitted the Journal of Marine Systems.

Ozgokmen, T. M., A. J. Mariano, L. I. Piterbarg, and A. Griffa , 1999: Predictability analysis of drifter clusters in the tropical Pacific Ocean. In preparation.